

Selberg Type Integrals Associated with $\mathfrak{sl}_3$

V.Tarasov^{*} and A.Varchenko^{*}

^{*}*St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg 191011, Russia*

^{*}*Department of Mathematics, University of North Carolina
Chapel Hill, NC 27599, USA*

February 2003

Abstract. We present several formulae for Selberg type integrals associated with the Lie algebra $\mathfrak{sl}_3$.

1. Introduction

The Selberg integral is the integral

$$(1.1) \quad \int_{\Delta^k[0,1]} \prod_{a=1}^k t_a^{\alpha-1} (1-t_a)^{\beta-1} \prod_{1 \leq a < b \leq k} (t_a - t_b)^{2\gamma} d^k t$$

where $\Delta^k[x, y] = \{t \in \mathbb{R}^k \mid x \leq t_k \leq \dots \leq t_1 \leq y\}$. Integral (1.1) is a generalization of the Euler beta integral. In 1944 Selberg showed [S] that the integral equals

$$\prod_{j=0}^{k-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\beta + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\alpha + \beta + (2k - 2 - j)\gamma) \Gamma(\gamma)}.$$

The Selberg integral is one of the most remarkable hypergeometric functions with many applications, see for instance [A1], [A2], [As], [D], [DF1], [DF2], [FSV], [M]. Taking a suitable limit of the Selberg integral one gets the exponential Selberg integral:

$$(1.2) \quad \int_{\Delta^k[0,+\infty]} \prod_{a=1}^k e^{-t_a} t_a^{\alpha-1} \prod_{1 \leq a < b \leq k} (t_a - t_b)^{2\gamma} d^k t = \prod_{j=0}^{k-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)}.$$

^{*}Supported in part by RFFI grant 02-01-00085a and CRDF grant RM1-2334-MO-02
E-mail: vt@pdmi.ras.ru

^{*}Supported in part by NSF grant DMS-9801582
E-mail: anv@email.unc.edu

There is also a discrete version of the exponential Selberg integral:

$$(1.3) \quad \sum_{u \in \Delta_{\mathbb{Z}, \gamma}^k} \prod_{a=1}^k z^{u_a} \frac{\Gamma(u_a + \alpha)}{\Gamma(u_a + 1)} \prod_{1 \leq a < b \leq k} \frac{(u_a - u_b) \Gamma(u_a - u_b + \gamma)}{\Gamma(u_a - u_b - \gamma + 1)} =$$

$$= z^{k(k-1)\gamma/2} (1 - z)^{-k\alpha - k(k-1)\gamma} \prod_{j=0}^{k-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)}$$

where $\Delta_{\mathbb{Z}, \gamma}^k = (k-1, k-2, \dots, 0)\gamma + (\Delta^k[0, +\infty] \cap \mathbb{Z}^k)$. Formula (1.3) can be obtained from the formula for the q -Selberg integral, see [AK], by the so-called rational degeneration.

The series (1.3) converges for $|z| < 1$ and any α, γ such that all gamma-functions are well defined. As $z \rightarrow 1$, formula (1.3) reproduces formula (1.2).

For generic α, γ one can replace the sum over the lattice cone $\Delta_{\mathbb{Z}, \gamma}^k$ by the sum over the total lattice

$$(1.4) \quad \mathbb{Z}_{\gamma}^k = (k-1, k-2, \dots, 0)\gamma + \mathbb{Z}^k,$$

since all additional terms appearing in the sum over $\mathbb{Z}_{\gamma}^k$ vanish.

All three types of Selberg integrals are related to representation theory of the Lie algebra $\mathfrak{sl}_2$. Namely, given a tensor product $V = V_1 \otimes \dots \otimes V_n$ of $\mathfrak{sl}_2$ -modules one has various linear systems of differential and difference equations for a V -valued function $u(z_1, \dots, z_n)$. The equations are called the Knizhnik-Zamolodchikov (KZ) and dynamical equations, see [EFK], [EV], [FMTV], [KZ], [TV2], [TV3], [V]. Explicit formulae can be found in [TV3]. All three types of Selberg integrals appear as coordinate functions of hypergeometric solutions of those equations, see [FMTV], [MaV], [TV1], [SV], [V].

In particular, integral (1.1) appears as a coordinate function of the hypergeometric solution of the KZ differential equations with values in the space

$$\text{Sing}(L_{\ell_1} \otimes L_{\ell_2})[\ell_1 + \ell_2 - 2k] = \{v \in L_{\ell_1} \otimes L_{\ell_2} \mid hv = (\ell_1 + \ell_2 - 2k)v, \quad ev = 0\},$$

where L_{ℓ} is the irreducible $\mathfrak{sl}_2$ -module with highest weight $\ell \in \mathbb{C}$, and e, h are the standard generators of the Borel subalgebra of $\mathfrak{sl}_2$.

Integral (1.2) appears as the coordinate function of the hypergeometric solution of the dynamical differential equation from [FMTV] with values in the weight subspace

$$L_{\ell}[\ell - 2k] = \{v \in L_{\ell} \mid hv = (\ell - 2k)v\}.$$

Integral (1.3) appears as the coordinate function of the hypergeometric solution of the dynamical differential equation from [TV3] with values in $L_{\ell}[\ell - 2k]$.

In all three cases the unknown function $u(z_1, \dots, z_n)$ takes values in a one-dimensional space. It was conjectured in [MV] that if the space of values of a KZ or a dynamical equation is one-dimensional, then coordinates of the hypergeometric solution can be expressed in terms of elementary functions and gamma-functions.

Motivated by this ideology, we give in this paper four versions of new Selberg type integrals analogous to integrals (1.1)–(1.3), see formulae (2.5), (3.2)–(3.4). The new integrals

are coordinate functions of hypergeometric solutions of KZ and dynamical equations associated with the Lie algebra $\mathfrak{sl}_3$, but we do not discuss this connection in the present paper. The consideration extends in a rather straightforward way to the $\mathfrak{sl}_n$ case for $n > 3$ giving the corresponding Selberg type integrals. This will be done in a separate paper.

The generalization of the Selberg integral proposed in this paper is complementary to the definition of Selberg type integrals associated with root systems [Ma]. In that definition the root system enters the integrand via the product of linear functions of integration variables, the linear functions defining the reflection hyperplanes of the root system. From that point of view the Selberg integral (1.1) is associated with the root system A_{k-1} . In our generalization of the Selberg integral the linear factors of the integrand do not depend on the root system, but the root system governs the exponents of linear factors of the integrand, cf. (3.3), (3.4). From our point of view the Selberg integral, defined for any k by formula (1.1), corresponds to the root system A_1 .

One can expect that there are Selberg type integrals depending on two root systems, the first one determining the linear factors of the integrand, like in [Ma], and the second one governing the exponents, like in this paper.

2. Discrete exponential Selberg integrals associated with $\mathfrak{sl}_3$

For any function $f(t_1, \dots, t_k)$ set

$$\text{Sym}_{t_1, \dots, t_k} f(t_1, \dots, t_k) = \frac{1}{k!} \sum_{\sigma \in \mathbf{S}_k} f(t_{\sigma_1}, \dots, t_{\sigma_k}).$$

Fix nonnegative integers k_1, k_2 such that $k_1 \geq k_2$. Introduce the master function

$$\begin{aligned} (2.1) \quad \Phi(u_1, \dots, u_{k_1}; v_1, \dots, v_{k_2}; z_1, z_2) = \\ = \prod_{a=1}^{k_1} z_1^{u_a} \frac{\Gamma(u_a + \alpha)}{\Gamma(u_a + 1)} \prod_{b=1}^{k_2} z_2^{v_b} \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} \frac{\Gamma(v_b - u_a - \gamma + 1)}{\Gamma(v_b - u_a + 1)} \times \\ \times \prod_{1 \leq a < b \leq k_1} \frac{(u_a - u_b) \Gamma(u_a - u_b + \gamma)}{\Gamma(u_a - u_b - \gamma + 1)} \prod_{1 \leq a < b \leq k_2} \frac{(v_a - v_b) \Gamma(v_a - v_b + \gamma)}{\Gamma(v_a - v_b - \gamma + 1)}, \end{aligned}$$

and the weight function

$$\begin{aligned} (2.2) \quad w(u_1, \dots, u_{k_1}; v_1, \dots, v_{k_2}) = \\ = \text{Sym}_{u_1, \dots, u_{k_1}} \text{Sym}_{v_1, \dots, v_{k_2}} \left(\prod_{b=1}^{k_2} \frac{1}{v_b - u_{b+k_1-k_2} - \gamma} \prod_{1 \leq b < a \leq k_2} \frac{v_b - u_{a+k_1-k_2}}{v_b - u_{a+k_1-k_2} - \gamma} \times \right. \\ \left. \times \prod_{1 \leq a < b \leq k_1} \frac{u_a - u_b - \gamma}{u_a - u_b} \prod_{1 \leq a < b \leq k_2} \frac{v_a - v_b - \gamma}{v_a - v_b} \right). \end{aligned}$$

The weight function is symmetric in $u_1, \dots, u_{k_1}$ and in $v_1, \dots, v_{k_2}$ separately, and has at

most simple poles located at the hyperplanes $v_b - u_a = \gamma$. Its numerator

$$w(u_1, \dots, u_{k_1}; v_1, \dots, v_{k_2}) \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} (v_b - u_a - \gamma)$$

vanishes at every triple intersection of hyperplanes of the form $u_a = u_b + \gamma = v_c$ or of the form $u_a = v_b = v_c - \gamma$. Set

$$(2.3) \quad \begin{aligned} F(u_1, \dots, u_{k_1}; v_1, \dots, v_{k_2}; z_1, z_2) &= \\ &= \Phi(u_1, \dots, u_{k_1}; v_1, \dots, v_{k_2}; z_1, z_2) w(u_1, \dots, u_{k_1}; v_1, \dots, v_{k_2}). \end{aligned}$$

Lemma 2.1. *Let $(\bar{u}, \bar{v}) \in \mathbb{Z}_\gamma^{k_1} \times \mathbb{Z}_\gamma^{k_2}$, cf. (1.4). Let α, γ be generic. Then the function $F(u; v; z_1, z_2)$ has a limit as $(u, v) \rightarrow (\bar{u}, \bar{v})$ along any straight line which does not belong to the singularity hyperplanes of the function F , and the limit does not depend on the direction. By abuse of notation we denote this limit $F(\bar{u}; \bar{v}; z_1, z_2)$.*

Actually, in a neighbourhood of any point $(\bar{u}, \bar{v}) \in \mathbb{Z}_\gamma^{k_1} \times \mathbb{Z}_\gamma^{k_2}$ the function F can be written in the form $F = G + P/Q$, where G, P, Q are functions, regular at $(\bar{u}, \bar{v})$, and the leading part of the Taylor series expansion of the function P at $(\bar{u}, \bar{v})$ has greater homogeneous degree than that of the function Q . Hence, the limit $F(\bar{u}; \bar{v}; z_1, z_2)$ equals $G(\bar{u}; \bar{v})$.

Let $\Delta^{k_1, k_2}[x, y]$ be a domain in $\mathbb{R}^{k_1+k_2}$ with coordinates $t_1, \dots, t_{k_1}, s_1, \dots, s_{k_2}$ defined by the following inequalities:

$$(2.4) \quad \begin{aligned} x \leq t_{k_1} \leq t_{k_1-1} \leq \dots \leq t_{k_1-k_2+1} \leq t_{k_1-k_2} \leq \dots \leq t_1 \leq y \\ \wedge \quad \quad \quad \wedge \quad \quad \quad \dots \quad \quad \quad \wedge \\ s_{k_2} \leq s_{k_2-1} \leq \dots \leq s_1 \leq y \end{aligned}.$$

Let $\Delta_{\mathbb{Z}, \gamma}^{k_1, k_2} = (k_1 - 1, \dots, 0, k_2 - 1, \dots, 0)\gamma + (\Delta^{k_1, k_2}[0, +\infty] \cap \mathbb{Z}^{k_1+k_2})$.

Lemma 2.2. *Let $(u, v) \in \mathbb{Z}_\gamma^{k_1} \times \mathbb{Z}_\gamma^{k_2}$. Let α, γ be generic. Then $F(u; v; z_1, z_2) = 0$ unless $(u, v) \in \Delta_{\mathbb{Z}, \gamma}^{k_1, k_2}$.*

Theorem 2.3. (Discrete exponential Selberg integral in the $\mathfrak{sl}_3$ case)

$$(2.5) \quad \begin{aligned} \sum_{(u, v) \in \Delta_{\mathbb{Z}, \gamma}^{k_1, k_2}} F(u; v; z_1, z_2) &= z_1^{k_1(k_1-1)\gamma/2} z_2^{k_2(k_2-1)\gamma/2} \times \\ &\times (1 - z_1)^{(k_1-k_2)(\gamma-\alpha-k_1\gamma)} (1 - z_2)^{k_2(k_1-k_2+1)\gamma} (1 - z_1 z_2)^{k_2(\gamma-\alpha-k_1\gamma)} \times \\ &\times \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_2-1} \frac{\Gamma(-k_1\gamma + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \end{aligned}$$

where the function F is defined by formulae (2.1)–(2.3).

The series (2.5) converges for $|z_1| < 1$, $|z_2| < 1$ and generic α, γ . For $k_2 = 0$ formula (2.5) coincides with formula (1.3).

3. Selberg integrals associated with $\mathfrak{sl}_3$

For any nondecreasing map $M : \{1, \dots, k_2\} \rightarrow \{1, \dots, k_1\}$ such that $M(b) \leq k_1 - k_2 + b$ for any $b = 1, \dots, k_2$, introduce a domain $D_M^{k_1, k_2}[x, y]$ in $\mathbb{R}^{k_1 + k_2}$ with coordinates $t_1, \dots, t_{k_1}, s_1, \dots, s_{k_2}$ defined by the following inequalities:

$$\begin{aligned} x \leq t_{k_1} \leq \dots \leq t_1 \leq y, \quad x \leq s_{k_2} \leq \dots \leq s_1 \leq y, \\ t_{M(b)} \leq s_b \leq t_{M(b)-1}, \quad b = 1, \dots, k_2. \end{aligned}$$

Here $t_0 = y$. Note that $\Delta^{k_1, k_2}[x, y] = \sum_M D_M^{k_1, k_2}[x, y]$ as chains. Consider the chain

$$C_\gamma^{k_1, k_2}[x, y] = \sum_M X_{M, \gamma}^{k_1, k_2} D_M^{k_1, k_2}[x, y]$$

with coefficients

$$X_{M, \gamma}^{k_1, k_2} = \prod_{b=1}^{k_2} \frac{\sin(\pi(k_1 - k_2 - M(b) + b + 1)\gamma)}{\sin(\pi(k_1 - k_2 + b)\gamma)}.$$

Introduce the weight function

$$\begin{aligned} (3.1) \quad g(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) &= \text{Sym}_{t_1, \dots, t_{k_1}} \text{Sym}_{s_1, \dots, s_{k_2}} \left(\prod_{b=1}^{k_2} \frac{1}{s_b - t_{b+k_1-k_2}} \right) = \\ &= \text{Sym}_{t_1, \dots, t_{k_1}} \text{Sym}_{s_1, \dots, s_{k_2}} \left(\prod_{b=1}^{k_2} \frac{1}{s_b - t_b} \right). \end{aligned}$$

The weight function is symmetric in $t_1, \dots, t_{k_1}$ and in $s_1, \dots, s_{k_2}$ separately, and has at most simple poles located at the hyperplanes $t_a = s_b$. Formula (3.1) is the specialization of formula (2.2) at $\gamma = 0$.

Theorem 3.1. (Exponential Selberg integral in the $\mathfrak{sl}_3$ case)

$$\begin{aligned} (3.2) \quad & \int_{C_\gamma^{k_1, k_2}[0, +\infty]} \prod_{a=1}^{k_1} e^{-\beta_1 t_a} t_a^{\alpha-1} \prod_{b=1}^{k_2} e^{-\beta_2 s_b} \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} |t_a - s_b|^{-\gamma} \times \\ & \times \prod_{1 \leq a < b \leq k_1} (t_a - t_b)^{2\gamma} \prod_{1 \leq a < b \leq k_2} (s_a - s_b)^{2\gamma} g(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) d^{k_1} t d^{k_2} s = \\ & = \beta_1^{(k_1 - k_2)(\gamma - \alpha - k_1 \gamma)} \beta_2^{k_2(k_1 - k_2 + 1)\gamma} (\beta_1 + \beta_2)^{k_2(\gamma - \alpha - k_1 \gamma)} \times \\ & \times \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_2-1} \frac{\Gamma(-k_1 \gamma + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)}. \end{aligned}$$

Theorem 3.2. (Selberg integral in the $\mathfrak{sl}_3$ case)

$$\begin{aligned}
(3.3) \quad & \int_{C_\gamma^{k_1, k_2}[0,1]} \prod_{a=1}^{k_1} t_a^{\alpha-1} (1-t_a)^{\beta_1-1} \prod_{b=1}^{k_2} (1-s_b)^{\beta_2-1} \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} |t_a - s_b|^{-\gamma} \times \\
& \times \prod_{1 \leq a < b \leq k_1} (t_a - t_b)^{2\gamma} \prod_{1 \leq a < b \leq k_2} (s_a - s_b)^{2\gamma} g(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) d^{k_1} t d^{k_2} s = \\
& = \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_1-k_2-1} \frac{\Gamma(\beta_1 + j\gamma)}{\Gamma(\alpha + \beta_1 + (2k_1 - k_2 - 2 - j)\gamma)} \times \\
& \times \prod_{j=0}^{k_2-1} \frac{\Gamma(\beta_2 + j\gamma) \Gamma(\beta_1 + \beta_2 - 1 - \gamma + j\gamma) \Gamma(-k_1\gamma + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\beta_2 + (2k_2 - k_1 - 2 - j)\gamma) \Gamma(\alpha + \beta_1 + \beta_2 - 1 + (k_1 + k_2 - 3 - j)\gamma) \Gamma(\gamma)} .
\end{aligned}$$


Theorem 3.3. (Another Selberg integral in the $\mathfrak{sl}_3$ case)

$$\begin{aligned}
(3.4) \quad & \int_{C_\gamma^{k_1, k_2}[0,1]} \prod_{a=1}^{k_1} t_a^{\alpha-1} (1-t_a)^{\beta_1-1} \prod_{b=1}^{k_2} (1-s_b)^{\beta_2-1} \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} |t_a - s_b|^{-\gamma} \times \\
& \times \prod_{1 \leq a < b \leq k_1} (t_a - t_b)^{2\gamma} \prod_{1 \leq a < b \leq k_2} (s_a - s_b)^{2\gamma} d^{k_1} t d^{k_2} s = \\
& = \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_1-k_2-1} \frac{\Gamma(\beta_1 + j\gamma)}{\Gamma(\alpha + \beta_1 + (2k_1 - k_2 - 2 - j)\gamma)} \times \\
& \times \prod_{j=0}^{k_2-1} \frac{\Gamma(\beta_2 + j\gamma) \Gamma(\beta_1 + \beta_2 - \gamma + j\gamma) \Gamma(1 - k_1\gamma + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\beta_2 + 1 + (2k_2 - k_1 - 2 - j)\gamma) \Gamma(\alpha + \beta_1 + \beta_2 + (k_1 + k_2 - 3 - j)\gamma) \Gamma(\gamma)} .
\end{aligned}$$

Integrals (3.2)–(3.4) converge if $\operatorname{Re} \alpha > 0$, $\operatorname{Re} \beta_1 > 0$, $\operatorname{Re} \beta_2 > 0$, $\operatorname{Re} \gamma < 0$ and $|\operatorname{Re} \gamma|$ is sufficiently small. For $k_2 = 0$ formulae (3.2) and (3.3), (3.4) coincide respectively with formulae (1.2) and (1.1).

There is another description of the integration chains $C_\gamma^{k_1, k_2}$ in Theorems 3.1–3.3. We explain it for the exponential Selberg integral (3.2). For the Selberg integrals (3.3) and (3.4) the description is similar.

Consider a collection of simple curves $\mathcal{C}_1, \dots, \mathcal{C}_{k_1+k_2}$ going around the real positive semi-line $\mathbb{R}_{\geq 0}$ like the curve in the picture:



and such that $\mathcal{C}_i$ is between $\mathbb{R}_{\geq 0}$ and $\mathcal{C}_j$ for any $i < j$. Let $G(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2})$ be the integrand in the left hand side of formula (3.2) and let

$$\begin{aligned} \tilde{G}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) &= \prod_{a=1}^{k_1} e^{-\beta_1 t_a} t_a^{\alpha-1} \prod_{b=1}^{k_2} e^{-\beta_2 s_b} \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} (s_b - t_a)^{-\gamma} \times \\ &\times \prod_{1 \leq a < b \leq k_1} (t_a - t_b)^{2\gamma} \prod_{1 \leq a < b \leq k_2} (s_a - s_b)^{2\gamma} g(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) \end{aligned}$$

(no absolute value of $s_b - t_a$). Fix a branch of the function $\tilde{G}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2})$ on $\mathcal{C}_1 \times \dots \times \mathcal{C}_{k_1+k_2}$ in the following way: at the point, where all the variables $t_1, \dots, t_{k_1}, s_1, \dots, s_{k_2}$ are real negative, one has $\arg t_a = \arg(s_b - t_a) = \pi$ for any a, b , and $\arg(t_a - t_b) = \arg(s_a - s_b) = 0$ for $a < b$. Equivalently, if $\operatorname{Im} t_a > 0$, $\operatorname{Im} s_b > 0$ for any a, b , and $0 < \operatorname{Re} t_{k_1} < \dots < \operatorname{Re} t_1 < \operatorname{Re} s_{k_2} < \dots < \operatorname{Re} s_1$, then $\arg t_a$, $\arg(s_b - t_a)$ for any a, b , and $\arg(t_a - t_b)$, $\arg(s_a - s_b)$ for $a < b$ are between $-\pi/2$ and $\pi/2$.

Theorem 3.4.

$$\begin{aligned} \int_{\mathcal{C}_1 \times \dots \times \mathcal{C}_{k_1+k_2}} \tilde{G}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) d^{k_1}t d^{k_2}s &= \\ = N_{k_1}(\alpha, \gamma) N_{k_2}(-k_1\gamma, \gamma) \int_{C_\gamma^{k_1, k_2}[0, +\infty]} G(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) d^{k_1}t d^{k_2}s. \end{aligned}$$

where

$$N_k(\alpha, \gamma) = \prod_{j=0}^{k-1} \frac{2i e^{\pi i \alpha} \sin(\pi(\alpha + j\gamma)) \sin(\pi(\gamma + j\gamma))}{\sin(\pi\gamma)}.$$

The proof is straightforward.

4. On proofs of Theorems 2.3, 3.1 – 3.3

All four types of new Selberg integrals (2.5), (3.2)–(3.4) are related to representation theory of the Lie algebra $\mathfrak{sl}_3$. They appear as coordinate functions of hypergeometric solutions of the KZ and dynamical equations associated with $\mathfrak{sl}_3$.

Let α_1, α_2 be simple roots of $\mathfrak{sl}_3$, and let ω_1, ω_2 be the fundamental weights. Denote by L_λ the irreducible $\mathfrak{sl}_3$ -module of highest weight λ . Series (2.5) appears as the coordinate function of the hypergeometric solution of the dynamical differential equations from [TV3] with values in the weight subspace $L_\lambda[\lambda - k_1\alpha_1 - k_2\alpha_2]$ for $\lambda \in \mathbb{C}\omega_1$, and integral (3.2) appears as the coordinate function of the hypergeometric solution of the dynamical differential equations from [FMTV] with values in the same weight subspace, see [FMTV, formula (8)]. Integrals (3.3), (3.4) appear as coordinate functions of the hypergeometric solution of the KZ differential equations with values in the space $\operatorname{Sing}(L_{\lambda_1} \otimes L_{\lambda_2})[\lambda_1 + \lambda_2 - k_1\alpha_1 - k_2\alpha_2]$ of singular weight vectors for $\lambda_1 \in \mathbb{C}\omega_1$, see [SV], and Chapter 12 in [V].

To prove Theorem 2.3 one considers expressions in both sides of formula (2.5) as functions of the variables z_1, z_2 . The dynamical differential equations in the case in question take the form

$$\begin{aligned}\frac{\partial \Psi}{\partial z_1} &= \left(\frac{k_1(k_1-1)\gamma}{2z_1} + \frac{(k_1-k_2)(\alpha-\gamma+k_1\gamma)}{1-z_1} + \frac{z_2 k_2(\alpha-\gamma+k_1\gamma)}{1-z_1 z_2} \right) \Psi, \\ \frac{\partial \Psi}{\partial z_2} &= \left(\frac{k_2(k_2-1)\gamma}{2z_1} - \frac{k_2(k_1-k_2+1)\gamma}{1-z_2} + \frac{z_1 k_2(\alpha-\gamma+k_1\gamma)}{1-z_1 z_2} \right) \Psi.\end{aligned}$$

The series in the left hand side solves this system and, hence, equals the product in the right hand side up to a factor which does not depend on z_1, z_2 . Since the values of both expressions coincide at $z_1 = z_2 = 0$, the proportionality factor equals 1, which completes the proof.

Theorem 3.1 is derived from Theorem 2.3 by taking the limit $z_1 \rightarrow 1, z_2 \rightarrow 1$. Namely, set $z_1 = e^{-\varepsilon\beta_1}, z_2 = e^{-\varepsilon\beta_2}, u_a = t_a/\varepsilon, v_b = s_b/\varepsilon$, and take the limit $\varepsilon \rightarrow 0$. If $\operatorname{Re} \alpha > 0, \operatorname{Re} \beta_1 > 0, \operatorname{Re} \beta_2 > 0, \operatorname{Re} \gamma < 0$ and $|\operatorname{Re} \gamma|$ is sufficiently small, then using a corollary of the Stirling formula:

$$\frac{\Gamma(x+c)}{\Gamma(x+d)} = x^{c-d}(1+o(1)), \quad \operatorname{Re} x \rightarrow +\infty,$$

one shows that formula (2.5) tends to formula (3.2) in the limit. This is quite similar to getting formula (1.2) from formula (1.3) in the limit $z \rightarrow 1$.

To obtain Theorem 3.2 one shows that expressions in both sides of formula (2.5) obey the same system of difference equations with respect to the shifts of the variables $\beta_1 \rightarrow \beta_1+1$ and $\beta_2 \rightarrow \beta_2+1$, see [MaV]; those difference equations are the dynamical difference equations from [TV2]. Therefore, those expressions are proportional up to a periodic function of β_1, β_2 . The periodic function can be found by comparing asymptotics as $\operatorname{Re} \beta_1 \rightarrow +\infty, \operatorname{Re} \beta_2 \rightarrow +\infty$. In this limit formula (3.3) reduces to formula (3.2), which shows that the periodic function is the constant equal to 1.

The proof of Theorem 3.3 is based on formula (3.3). It is outlined in the next section, cf. (5.5).

5. Further generalizations of Selberg type integrals

Denote by $S(\alpha, \beta)$ the Selberg integral (1.1). Consider the following integrals:

$$I_l = \int_{\Delta^k[0,1]} \prod_{a=1}^k t_a^{\alpha-1} (1-t_a)^{\beta-1} \prod_{1 \leq a < b \leq k} (t_a - t_b)^{2\gamma} \operatorname{Sym}_{t_1, \dots, t_k} \left(\prod_{a=1}^l t_a \prod_{b=l+1}^k (1-t_a) \right) d^k t,$$

$l = 0, \dots, k$. In particular, $I_0 = S(\alpha, \beta+1)$ and $I_k = S(\alpha+1, \beta)$.

Proposition 5.1. *For any $l = 0, \dots, k$ one has*

$$I_l = \prod_{i=0}^{l-1} \frac{\alpha + (k-1-i)\gamma}{\beta + i\gamma} \prod_{j=0}^{k-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(1 + \beta + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(1 + \alpha + \beta + (2k-2-j)\gamma) \Gamma(\gamma)}.$$

The proof is straightforward using Aomoto's formula [A1]:

$$\begin{aligned} & \int_{\Delta^k[0,1]} \prod_{a=1}^k t_a^{\alpha-1} (1-t_a)^{\beta-1} \prod_{1 \leq a < b \leq k} (t_a - t_b)^{2\gamma} \operatorname{Sym}_{t_1, \dots, t_k} (t_1 \dots t_k) d^k t = \\ &= \prod_{i=0}^{l-1} \frac{\alpha + (k-1-i)\gamma}{\alpha + \beta + (2k-2-i)\gamma} \prod_{j=0}^{k-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\beta + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\alpha + \beta + (2k-2-j)\gamma) \Gamma(\gamma)}. \end{aligned}$$

Proposition 5.1 is equivalent to the formula for the Selberg integral and linear relations for the integrals $I_0, \dots, I_k$:

$$(\alpha + (k-l-1)\gamma) I_l = (\beta + l\gamma) I_{l+1}, \quad l = 0, \dots, k-1.$$

These relations correspond to the fact that the vector $\sum_{a=0}^k (-1)^a I_a f^{k-a} v_1 \otimes f^a v_2$ is a singular vector in the tensor product $L_{\ell_1} \otimes L_{\ell_2}$ of highest weight $\mathfrak{sl}_2$ -modules with highest weights $\ell_1 = -\alpha/\gamma$, $\ell_2 = -\beta/\gamma$, see [V]. Recall that a vector v is called singular if $ev = 0$. Here e, f are the standard generators of the opposite nilpotent subalgebras of the Lie algebra $\mathfrak{sl}_2$, and v_1, v_2 are the highest weight vectors (standard cyclic vectors) of the respective $\mathfrak{sl}_2$ -modules L_{ℓ_1}, L_{ℓ_2} .

Fix nonnegative integers k_1, k_2 such that $k_1 \geq k_2$. Denote by $R(\alpha, \beta_1, \beta_2)$ the integral (3.3). Consider the master function

$$\begin{aligned} & \Omega(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}; \alpha, \beta_1, \beta_2) = \\ &= \prod_{a=1}^{k_1} t_a^{\alpha-1} (1-t_a)^{\beta_1-1} \prod_{b=1}^{k_2} (1-s_b)^{\beta_2-1} \prod_{a=1}^{k_1} \prod_{b=1}^{k_2} |t_a - s_b|^{-\gamma} \times \\ & \quad \times \prod_{1 \leq a < b \leq k_1} (t_a - t_b)^{2\gamma} \prod_{1 \leq a < b \leq k_2} (s_a - s_b)^{2\gamma}. \end{aligned}$$

Say that a triple of nonnegative integers (l_1, l_2, m) is admissible if $l_1 \leq k_1 - k_2 + l_2$, $l_2 \leq k_2$ and $m \leq \min(l_1, l_2)$. For any admissible triple l_1, l_2, m introduce the functions

$$\begin{aligned} & h_{l_1, l_2, m}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) = \\ &= \operatorname{Sym}_{t_1, \dots, t_{k_1}} \operatorname{Sym}_{s_1, \dots, s_{k_2}} \left(\prod_{a=1}^{l_1} t_a \prod_{a=l_1+1}^{k_1} (1-t_a) \prod_{b=1}^m \frac{1-s_b}{s_b - t_b} \prod_{b=l_2+1}^{k_2} \frac{1-s_b}{s_b - t_{b+k_1-k_2}} \right), \\ & \tilde{h}_{l_1, l_2, m}(t_1, \dots, t_{k_1}; s_1, \dots, s_{k_2}) = \\ &= \operatorname{Sym}_{t_1, \dots, t_{k_1}} \operatorname{Sym}_{s_1, \dots, s_{k_2}} \left(\prod_{a=1}^{l_1} t_a \prod_{a=l_1+1}^{k_1} (1-t_a) \prod_{b=1}^m \frac{1-t_b}{s_b - t_b} \prod_{b=l_2+1}^{k_2} \frac{1-s_b}{s_b - t_{b+k_1-k_2}} \right), \end{aligned}$$

and the integrals

$$J_{l_1, l_2, m} = \int_{C_\gamma^{k_1, k_2}[0,1]} \Omega(t; s; \alpha, \beta_1, \beta_2) h_{l_1, l_2, m}(t; s) d^{k_1}t d^{k_2}s,$$

$$\tilde{J}_{l_1, l_2, m} = \int_{C_\gamma^{k_1, k_2}[0,1]} \Omega(t; s; \alpha, \beta_1, \beta_2) \tilde{h}_{l_1, l_2, m}(t; s) d^{k_1}t d^{k_2}s.$$

In particular, $J_{0,0,0} = \tilde{J}_{0,0,0} = R(\alpha, \beta_1 + 1, \beta_2 + 1)$. One can also observe that

$$(5.1) \quad J_{0,l,0}(\alpha + 1, \beta_1, \beta_2) = J_{k_1, k_2, k_2-l}(\alpha, \beta_1 + 1, \beta_2).$$

Here we indicate the dependence on the parameters α, β_1, β_2 explicitly. In addition, the integral

$$(5.2) \quad J_{0,k_2,0} = \tilde{J}_{0,k_2,0} = \int_{C_\gamma^{k_1, k_2}[0,1]} \Omega(t; s; \alpha, \beta_1 + 1, \beta_2) d^{k_1}t d^{k_2}s$$

is expressed in terms of the master function only with no extra rational functions involved.

Theorem 5.2. *Let (l_1, l_2, m) be an admissible triple. If $l_1 < k_1 - k_2 + l_2$, then*

$$\begin{aligned} & (\alpha + (k_1 - l_1 - 1)\gamma) J_{l_1, l_2, m} = \\ & = (\beta_1 + (l_1 - l_2 + m)\gamma) J_{l_1+1, l_2, m} - (l_2 - m)\gamma J_{l_1+1, l_2, m+1}, \end{aligned}$$

and if $m < l_2 < k_2$, then

$$\begin{aligned} (5.3) \quad & (k_2 - k_1 + l_1 - l_2)\gamma J_{l_1, l_2-1, m} = \\ & = (\beta_2 + (l_2 - m - 1)\gamma) J_{l_1, l_2, m} + (l_1 - m)\gamma J_{l_1, l_2, m+1}. \end{aligned}$$

Similarly, if $l_1 < k_1 - k_2 + l_2$, then

$$\begin{aligned} & (\alpha + (k_1 - l_1 - 1)\gamma) \tilde{J}_{l_1, l_2, m} = \\ & = (\beta_1 + (l_1 - m)\gamma) \tilde{J}_{l_1+1, l_2, m} - (l_2 - m)\gamma \tilde{J}_{l_1+1, l_2, m+1}, \end{aligned}$$

and if $m < l_2 < k_2$, then

$$\begin{aligned} (5.4) \quad & (k_2 - k_1 + l_1 - l_2)\gamma \tilde{J}_{l_1, l_2-1, m} = \\ & = (\beta_2 + (l_2 - l_1 + m - 1)\gamma) \tilde{J}_{l_1, l_2, m} + (l_1 - m)\gamma \tilde{J}_{l_1, l_2, m+1}. \end{aligned}$$

In the equations of Theorem 5.2 some of the triples $(l_1 + 1, l_2, m + 1)$, $(l_1, l_2 - 1, m)$, $(l_1, l_2, m + 1)$ may be inadmissible. In that case the corresponding coefficient:

$$l_2 - m, \quad k_2 - k_1 + l_1 - l_2, \quad l_1 - m,$$

respectively, equals zero, and the integral with an inadmissible index does not enter the equation.

Let $f_1, f_2 \in \mathfrak{sl}_3$ be root vectors corresponding to simple roots, and let ω_1, ω_2 be the fundamental weights. It is known that the vectors

$$\sum_{l_1, l_2, m} (-1)^{l_1} J_{l_1, l_2, m} \frac{f_1^{k_1 - k_2 - l_1 + l_2} [f_1, f_2]^{k_2 - l_2} v_1 \otimes f_1^{l_1 - m} [f_1, f_2]^m f_2^{l_2 - m} v_2}{(k_1 - k_2 - l_1 + l_2)! (k_2 - l_2)! (l_1 - m)! m! (l_2 - m)!}$$

and

$$\sum_{l_1, l_2, m} (-1)^{l_1} \tilde{J}_{l_1, l_2, m} \frac{f_1^{k_1 - k_2 - l_1 + l_2} [f_1, f_2]^{k_2 - l_2} v_1 \otimes f_2^{l_2 - m} [f_1, f_2]^m f_1^{l_1 - m} v_2}{(k_1 - k_2 - l_1 + l_2)! (k_2 - l_2)! (l_1 - m)! m! (l_2 - m)!}$$

are singular vectors, that is, they are annihilated by the action of the opposite root vectors $e_1, e_2 \in \mathfrak{sl}_3$, in the tensor product $L_{\lambda_1} \otimes L_{\lambda_2}$ of $\mathfrak{sl}_3$ -modules with highest weights

$$\lambda_1 = -\omega_1 \alpha / \gamma, \quad \lambda_2 = -\omega_1 \beta_1 / \gamma - \omega_2 \beta_2 / \gamma$$

and highest weight vectors v_1, v_2 , see [V]. This fact implies the relations of Theorem 5.2.

For generic values of $\alpha, \beta_1, \beta_2, \gamma$ one can solve equations of Theorem 5.2 and express all the integrals $J_{l_1, l_2, m}$ and $\tilde{J}_{l_1, l_2, m}$ via the known integral

$$J_{0,0,0} = \tilde{J}_{0,0,0} = R(\alpha, \beta_1 + 1, \beta_2 + 1).$$

For instance, the integrals $J_{0,l,0} = \tilde{J}_{0,l,0}$ can be easily computed using relations (5.3):

$$J_{0,l,0} = (-1)^l \prod_{i=0}^{l-1} \frac{(k_1 - k_2 + 1 + i)\gamma}{\beta_2 + i\gamma} J_{0,0,0}.$$

In particular,

$$(5.5) \quad J_{0,k_2,0} = \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_1-k_2-1} \frac{\Gamma(\beta_1 + 1 + j\gamma)}{\Gamma(\alpha + \beta_1 + 1 + (2k_1 - k_2 - 2 - j)\gamma)} \times \prod_{j=0}^{k_2-1} \frac{\Gamma(\beta_2 + j\gamma) \Gamma(\beta_1 + \beta_2 + 1 - \gamma + j\gamma) \Gamma(1 - k_1\gamma + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\beta_2 + 1 + (2k_2 - k_1 - 2 - j)\gamma) \Gamma(\alpha + \beta_1 + \beta_2 + 1 + (k_1 + k_2 - 3 - j)\gamma) \Gamma(\gamma)}.$$

After replacement $\beta_1 \rightarrow \beta_1 - 1$ the last formula yields formula (3.4), cf. (5.2). Similarly, the integrals $\tilde{J}_{k_1, k_2, m}$ can be computed using relations (5.4) and the integral $J_{k_1, k_2, 0} = \tilde{J}_{k_1, k_2, 0}$:

$$\begin{aligned} \tilde{J}_{k_1, k_2, m} &= (-1)^m \prod_{i=0}^{m-1} \frac{\beta_2 + (k_2 - k_1 - 1 + i)\gamma}{(k_1 - i)\gamma} J_{k_1, k_2, 0}, \\ J_{k_1, k_2, 0} &= \prod_{j=0}^{k_1-1} \frac{\Gamma(\alpha + 1 + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\gamma)} \prod_{j=0}^{k_1-k_2-1} \frac{\Gamma(\beta_1 + j\gamma)}{\Gamma(\alpha + \beta_1 + 1 + (2k_1 - k_2 - 2 - j)\gamma)} \times \\ &\times \prod_{j=0}^{k_2-1} \frac{\Gamma(\beta_2 + j\gamma) \Gamma(\beta_1 + \beta_2 - \gamma + j\gamma) \Gamma(1 - k_1\gamma + j\gamma) \Gamma(\gamma + j\gamma)}{\Gamma(\beta_2 + 1 + (2k_2 - k_1 - 2 - j)\gamma) \Gamma(\alpha + \beta_1 + \beta_2 + 1 + (k_1 + k_2 - 3 - j)\gamma) \Gamma(\gamma)}, \end{aligned}$$

the last formula following from (5.1), (5.5).

Errata

The published version of the paper (Lett. Math. Phys. **65** (2003), no. 3, 173–185) contains a sign misprint in the formulae for the functions $h_{l_1, l_2, m}$ and $\tilde{h}_{l_1, l_2, m}$ in Section 5. This misprint is corrected in the present version.

We thank O. Warnaar for a valuable remark.

References

- [A1] K. Aomoto, *Jacobi polynomials associated with Selberg integrals*, SIAM J. Math. Anal. **18** (1987), 545–549.
- [A2] K. Aomoto, *On the complex Selberg integral*, Q. J. Math. Oxford **38** (1987), 385–399.
- [AK] K. Aomoto and Y. Kato, *Gauss decomposition of connection matrices for symmetric A-type Jackson integrals*, Selecta Math., New Series **1** (1995), no. 4, 623–666.
- [As] R. Askey, *Some basic hypergeometric extensions of integrals of Selberg and Andrews*, SIAM J. Math. Anal. **11** (1980), 938–951.
- [D] V. Dotsenko, *Solving the $SU(2)$ conformal field theory with the Wakimoto free-field representation*, (in Russian) (1990), 1–31.
- [DF1] V. Dotsenko and V. Fateev, *Conformal algebra and multi-point correlation functions in 2-D statistical models*, Nucl. Phys. B **240** (1984), 312–348.
- [DF2] V. Dotsenko and V. Fateev, *Four-point correlation functions and the operator algebra in 2-D conformal invariant theories with central charge $C \leq 1$* , Nucl. Phys. B **251** (1985), 691–734.
- [EFK] P. Etingof, I. Frenkel and A. Kirillov Jr., *Lectures on representation theory and Knizhnik-Zamolodchikov equations*, Mathematical Surveys and Monographs, vol. 58, AMS, Providence RI, 1998.
- [EV] P. Etingof and A. Varchenko, *Dynamical Weyl groups and applications*, Adv. Math. **167** (2002), no. 1, 74–127.
- [FMTV] G. Felder, Ya. Markov, V. Tarasov and A. Varchenko, *Differential equations compatible with KZ equations*, Math. Phys., Analysis and Geometry **3** (2000), 139–177.

- [FSV] G. Felder, L. Stevens and A. Varchenko, *Elliptic Selberg integrals and conformal blocks*, Preprint (2002), 1–13, [math.QA/0210040](#).
- [FV1] G. Felder and A. Varchenko, *Integral representations for solutions of the elliptic Knizhnik-Zamolodchikov-Bernard equation*, Int. Math. Res. Notices (1995), no. 5, 221–233.
- [KZ] V. Knizhnik and A. Zamolodchikov, *Current algebra and Wess-Zumino model in two dimensions*, Nucl. Phys. B **247** (1984), 83–103.
- [Ma] I. G. Macdonald, *Some conjectures for root systems*, SIAM J. Math. Anal. **13** (1982), 988–1007.
- [M] M. L. Mehta, *Random Matrices*, Academic Press, 1991.
- [MaV] Ya. Markov and A. Varchenko, *Solutions of trigonometric KZ equations satisfy dynamical difference equations*, Adv. Math. **166** (2002), no. 1, 100–147.
- [MV] E. Mukhin and A. Varchenko, *Remarks on critical points of phase functions and norms of Bethe vectors*, Adv. Studies in Pure Math. **27**, 2000, pp. 239–246.
- [S] A. Selberg, *Bemerkninger om et multipliet integral*, Norsk Mat. Tidsskr. **26** (1944), 71–78.
- [SV] V. V. Schechtman and A. N. Varchenko, *Arrangements of hyperplanes and Lie algebras homology*, Invent. Math. **106** (1991), 139–194.
- [TV1] V. Tarasov and A. Varchenko, *Geometry of q -hypergeometric functions as a bridge between Yangians and quantum affine algebras*, Invent. Math. **128** (1997), no. 3, 501–588.
- [TV2] V. Tarasov and A. Varchenko, *Difference equations compatible with trigonometric KZ differential equations*, Int. Math. Res. Notices (2000), no. 15, 801–829..
- [TV3] V. Tarasov and A. Varchenko, *Duality for Knizhnik-Zamolodchikov and dynamical equations*, Acta Appl. Math. **73** (2002), no. 1-2, 141–154.
- [V] A. Varchenko, *Multidimensional hypergeometric functions and representation theory of Lie algebras and quantum groups*, Advanced Series in Math. Phys., vol. 21, World Scientific, Singapore, 1995.